

Osh4p is needed to reduce the level of phosphatidylinositol-4-phosphate on secretory vesicles as they mature

Yading Ling, Scott Hayano and Peter Novick

Department of Cellular and Molecular Medicine

University of California, San Diego, CA 92093

Send correspondence to:

Peter Novick

Department of Cellular and Molecular Medicine

GPL 238A

UCSD

9500 Gilman Drive,

La Jolla, CA 92093

pnovick@ucsd.edu

858 246 0465

Abstract

Phosphatidylinositol-4-phosphate (PI4P) is produced on both the Golgi and the plasma membrane. Despite extensive vesicular traffic between these compartments, genetic analysis suggests that the two pools of PI4P do not efficiently mix with one another. Several lines of evidence indicate that the PI4P produced on the Golgi is normally incorporated into secretory vesicles, but the fate of that pool has been unclear. We show here that in yeast the oxysterol binding proteins Osh1-Osh7 are collectively needed to maintain the normal

distribution of PI4P and that Osh4p is critical in this function. Osh4p associates with secretory vesicles at least in part through its interaction with PI4P and is needed, together with lipid phosphatases, to reduce the level of PI4P as vesicles approach sites of exocytosis. This reduction in PI4P is necessary for a switch in the regulation of the Sec4p exchange protein, Sec2p, from an interaction with the upstream Rab, Ypt31/32 to an interaction with a downstream Sec4p effector, Sec15p. Spatial regulation of PI4P levels thereby plays an important role in vesicle maturation.

Introduction

The various organelles of the exocytic and endocytic pathways each have unique functions that are dictated by their unique compositions. These distinct identities are maintained despite high rates of bi-directional vesicular traffic along and between these pathways. Maintenance of organelle identity requires both tightly regulated cargo selection during vesicle budding and accurate vesicle targeting to the appropriate compartment prior to fusion. Peripherally associated coat proteins direct cargo selection, while targeting requires several different regulatory mechanisms that work in concert. SNARE complexes, involving integral membrane proteins on the vesicle and target membranes, provide one level of specificity to the membrane fusion reaction (McNew *et al.*, 2000), however, because vesicle SNAREs must recycle back to the donor compartment after fusion, they are necessarily found on both anterograde and retrograde vesicles and therefore cannot specify the direction of vesicular traffic. Molecular tags that can be removed from the membrane after fusion are better suited to this aspect of vesicle targeting (Behnia and Munro, 2005).

Rab GTPases act as transient tags since they are activated by guanine nucleotide exchange factors (GEFs) and inactivated by GTPase activating proteins (GAPs) (Hutagalung and Novick, 2011). By localizing a GEF on one compartment and a GAP on another compartment an asymmetric distribution of activated Rab protein can be established that provides directional information to vesicular traffic. Once activated, Rabs recruit a variety of effectors such as molecular motors, tethering complexes and regulators of SNARE complex assembly that together define where a vesicle is delivered, tethered and fused (Guo *et al.*, 1999b; Grosshans *et al.*, 2006; Jin *et al.*, 2011; Mizuno-Yamasaki *et al.*, 2012).

Phosphoinositides can also serve as transient tags in membrane traffic pathways. Phosphorylation of the inositol head group of phosphatidylinositol at different positions generates distinct tags. By localizing a lipid kinase on one compartment and the competing lipid phosphatase on another a gradient can be established that provides spatial information for vectorial traffic (Behnia and Munro, 2005; Balla, 2013). These two types of tags can be exploited in a combinatorial fashion by proteins that act as coincidence detectors, binding only to membrane domains carrying both the appropriate Rab and phosphoinositide (Carlton and Cullen, 2005).

The yeast exocytic pathway is regulated by a series of Rabs. Ypt1p controls ER to Golgi transport as well as transport through the Golgi (Bacon *et al.*, 1989; Jedd *et al.*, 1995; Sclafani *et al.*, 2010). Ypt31p and Ypt32p are redundant Rabs that control exit from the Golgi (Benli *et al.*, 1996; Jedd *et al.*,

1997). Sec4p controls the delivery of secretory vesicles along polarized actin cables, vesicle tethering by the exocyst complex to exocytic sites specified by polarity establishment machinery and SNARE-mediated fusion to the plasma membrane (Guo *et al.*, 1999b; Grosshans *et al.*, 2006; Shen *et al.*, 2013). Sec4p is activated by its GEF, Sec2p, which is in turn regulated by several different mechanisms (Walch-Solimena *et al.*, 1997). Sec2p is recruited to membranes by the combinatorial signals of Ypt31/32-GTP and the phosphoinositide, PI4P (Ortiz *et al.*, 2002; Mizuno-Yamasaki *et al.*, 2010). Sec2p also binds to Sec15p, a subunit of the exocyst complex and a direct downstream effector of Sec4-GTP (Guo *et al.*, 1999b; Medkova *et al.*, 2006). The interaction of Sec2p, a Sec4p GEF, with Sec15p, a Sec4p effector, could generate a positive feedback loop leading to formation of a domain of highly activated Sec4p and highly concentrated exocyst. The binding sites for Ypt31/32-GTP and Sec15p overlap such that Sec2p can only bind to one or the other (Medkova *et al.*, 2006). This choice is dictated in part by the local PI4P concentration since high levels of PI4P prevent binding of Sec15p to Sec2p, but do not interfere with binding to Ypt31/32 (Mizuno-Yamasaki *et al.*, 2010).

Two PI4 kinases generate most of the PI4P in yeast; Pik1p is found in the Golgi (Walch-Solimena and Novick, 1999) and generates the Golgi-associated pool of PI4P, while Stt4p is found on the plasma membrane and generates a similar amount of PI4P at this site (Audhya *et al.*, 2000; Audhya and Emr, 2002). Although Golgi-derived vesicles constantly deliver new membrane to the cell surface, the Golgi pool of PI4P is not efficiently transferred to the plasma

membrane as indicated by the observation that even overexpression of Pik1p fails to complement the loss of Stt4p (Audhya *et al.*, 2000). This finding suggests that the Golgi pool of PI4P either fails to be incorporated into secretory vesicles or the vesicular pool of PI4P is reduced by one of several possible mechanisms. Here we explore the role of the Osh proteins in the regulation of the secretory vesicle pool of PI4P.

Results

Osh proteins are key regulators of PI4P distribution in cells

The phosphoinositide PI4P is critical for the localization of a number of proteins involved in polarized secretion, including Sec2p (Mizuno-Yamasaki *et al.*, 2010). Here we investigate how the level of PI4P is regulated on secretory vesicles. Studies from several groups have demonstrated that the oxysterol binding proteins (Osh proteins) control the level and intracellular distribution of PI4P in the budding yeast *Saccharomyces cerevisiae* (LeBlanc and McMaster, 2010; Stefan *et al.*, 2011). To begin our analysis of the role of Osh proteins in PI4P distribution, we employed cells lacking all endogenous Osh proteins and solely expressing a temperature-sensitive *osh4^{ts}* allele. Consistent with a prior report (Stefan *et al.*, 2011), we observed that the PI4P probe, GFP-PH^{FAPP1}, consisting of the PH domain from the FAPP1 protein fused to GFP, shifted from the Golgi apparatus to the plasma membrane in the *osh1-7Δ osh4^{ts}* cells after a temperature shift to 37 °C (Figure 1A). In addition, there was significantly more GFP-PH^{FAPP1} at polarized secretory sites, such as the tips of small buds and the necks of large budded cells, in *osh1-7Δ osh4^{ts}* cells at the non-permissive temperature (Figure 1A). These GFP-PH^{FAPP1} positive structures could represent

either secretory vesicles containing abnormally high levels of PI4P or an accumulation of PI4P produced at the plasma membrane. To differentiate these two possibilities, we labeled the plasma membrane with the lipophilic dye FM4-64. We noticed that the GFP-PH^{FAPP1} positive structures accumulated in the small buds of *osh1-7Δ osh4^{ts}* cells were clearly internal to the FM4-64 labeled plasma membrane (Figure 1B). These results suggest that one or more Osh proteins are needed to reduce the level of PI4P on secretory vesicles.

To investigate whether PI4P is present on secretory vesicles in a wild type *OSH* background, we examined the localization of another PI4P probe, consisting of the PH domain from Osh2p fused to GFP (Roy and Levine, 2004), in a *sec6^{ts}* mutant allele. We used a second probe for PI4P since the PH domain of FAPP1 has affinity not only for PI4P, but also Arf1p and may preferentially label membranes carrying both ligands (Roy and Levine, 2004). Sec6 is an essential component of the exocyst complex that is required for tethering secretory vesicles to the plasma membrane, and loss of Sec6 function leads to the accumulation of secretory vesicles (TerBush *et al.*, 1996). Vesicle accumulation is initially confined to regions of active secretion and surface expansion, but after a prolonged shift to the restrictive temperature vesicles accumulate throughout the cell (Govindan *et al.*, 1995). If PI4P is present on secretory vesicles we would expect accumulation of the GFP-2xPH^{Osh2} PI4P probe at polarized growth sites in *sec6^{ts}* mutant cells after short incubations at the non-permissive temperature. At the permissive temperature, we noticed that GFP-2xPH^{Osh2} was present both on the plasma membrane and Golgi apparatus

as previously reported (Roy and Levine, 2004). Intriguingly, upon shifting the *sec6^{ts}* mutant cells to 37°C for only 15 minutes, a notable increase of GFP-2xPH^{Osh2} positive structures were observed at the necks of large budded cells (Figure 1C). We observed only a slight increase of GFP-2xPH^{Osh2} positive structures near the cell necks in wild type cells upon a temperature shift (Figure 1C). These results provide additional evidence that PI4P is present in the membrane of secretory vesicles.

Osh proteins are key regulators of secretion in yeast

PI4P is a major regulator of Golgi to PM vesicle transport and loss of Osh protein function affects the secretion of a number of cargo proteins including Bgl2 in yeast cells (Kozminski *et al.*, 2006). To examine whether Osh proteins affect the fusion of Golgi-derived vesicles to the plasma membrane, we examined *osh1-7Δ osh4^{ts}* cells and *osh1-7Δ OSH4* cells at 37°C using thin section electron microscopy. Consistent with prior reports (Beh and Rine, 2004), we found that there was a large increase in the number of vesicles present in the cytoplasm of the *osh1-7Δ osh4^{ts}* cells (Supplemental Figure 1A; 1.72 vesicles/ μm^2) compared to cells containing wild type *OSH4* (Supplemental Figure 1A; 0.25 vesicles/ μm^2) under this condition.

Osh proteins have been implicated in regulation of assembly of the exocyst, a protein complex required for tethering secretory vesicles to the plasma membrane (TerBush *et al.*, 1996; Alfaro *et al.*, 2011). We examined the localizations of two subunits of the exocyst complex in the *osh1-7Δ osh4^{ts}* cells, Sec15p, an exocyst subunit that associates with secretory vesicles as they are

delivered to sites of secretion, and Sec3p, an exocyst subunit that can directly recognize exocytic sites on the plasma membrane. Both proteins were mislocalized from polarized secretion sites in over 90% of the *osh1-7Δ osh4^{ts}* cells when they were shifted from the permissive temperature to the non-permissive temperature (Supplemental Figure 1B). Meanwhile, Sec15p and Sec3p still localized normally in the *osh1-7Δ OSH4* cells upon the temperature shift (Supplemental Figure 1B). Together, these results suggest that Osh protein function is key to the tethering and/or fusion of secretory vesicles *in vivo*, presumably by controlling the level and intracellular distribution of PI4P.

Osh4 localizes to Sec4-positive vesicles

There are seven different Osh proteins in the budding yeast *Saccharomyces cerevisiae*. They share redundant yet essential roles *in vivo* as any one is sufficient for cell growth but collectively they are required for cell viability (Beh *et al.*, 2001). Osh1p localizes to Golgi and nuclear-vacuolar junction sites (Levine and Munro, 2001), while Osh2p, Osh3p, Osh6p and Osh7p have been previously localized to the ER-PM contact sites (Levine and Munro, 2001; Schulz *et al.*, 2009; Stefan *et al.*, 2011). To investigate which Osh protein is the master regulator of PI4P on secretory vesicles, we tagged Osh4p and Osh5p with GFP, and examined their *in vivo* localization using fluorescence microscopy. Only Osh4-GFP displayed a good signal by fluorescence microscopy (our unpublished observations). To enhance the signal, we constructed a 3xGFP fusion at the C-terminus of Osh4p. We observed three major pools of Osh4-3xGFP present in wild type cells. One pool was present on punctate structures in cytoplasm, while

another pool was concentrated at sites of polarized secretion, including the tips of small buds and mother-daughter necks. However, the majority of Osh4-3xGFP was cytoplasmic (Figure 2A). The pool of Osh4-3xGFP found in small bud tips colocalized well with mCherry tagged Sec4 (Figure 2A), suggesting that it is present on secretory vesicles and therefore a good candidate for the major Osh protein regulating the final stage of the secretory pathway.

Secretory vesicles are normally delivered to sites of polarized growth by the type V myosin motor, Myo2p (Govindan *et al.*, 1995; Schott *et al.*, 1999; Schott *et al.*, 2002). Myo2p is recruited to vesicles by Sec4-GTP and therefore in mutants defective in Sec4p activation vesicles accumulate in a depolarized fashion (Walch-Solimena *et al.*, 1997; Wagner *et al.*, 2002; Donovan and Bretscher, 2012). To determine if Osh4p is associated with secretory vesicles, we examined Osh4-3xGFP localization in *sec4^{ts}* mutant cells. At the permissive temperatures Osh4-3xGFP was present, in part, at polarized secretion sites, including small bud tips and mother-daughter cell necks (Figure 2B). However, upon shifting the cells to the non-permissive temperature for 1 hour, the polarized localization of Osh4-3xGFP was greatly reduced and we observed some abnormal punctate structures at the cell perimeter (Figure 2B and Figure 2C). Osh4-3xGFP still localized normally in wild type cells under the same condition (Figure 2B and Figure 2C). To further investigate whether Sec4p controls Osh4p localization *in vivo*, we performed a subcellular fractionation experiment. In *sec4^{ts}* cells, roughly 83% of Osh4p was in the S100 fraction at the permissive temperature. However, upon shifting the cells to the non-permissive temperature

for 1 hour, the percentage of Osh4p in the soluble fraction increased; with 91% in the S100 fraction (Supplemental Figure 2A and 2B). These results suggest Sec4p contributes to Osh4p's association with secretory vesicles.

To test the possibility that Osh4p is recruited to secretory vesicles directly by Sec4p, we over-expressed Sec4p several fold *in vivo*. However, we did not observe a detectable increase of polarized Osh4-3xGFP (Supplemental Figure 3A). In addition, we tested whether Sec4p directly binds Osh4p. Although we cannot rule out that there might be some weak interaction between Sec4p and Osh4p *in vivo*, we did not observe a direct interaction of these two proteins *in vitro*, regardless of the nucleotide loading status of Sec4p (Supplemental Figure 3B). We also tested Osh4p's binding to the upstream rab, Ypt32p, yet did not observe any interaction (Supplemental Figure 3C). These results support the proposal that Osh4p is associated with secretory vesicles and, although Sec4p contributes to the recruitment of Osh4p, it is not the only factor or switch controlling Osh4p localization *in vivo*.

The polarized localization of Osh4 is dependent on binding to PI4P

Osh4p binds to both ergosterol and PI4P (Im *et al.*, 2005; de Saint-Jean *et al.*, 2011). A prior study demonstrated that ergosterol is important for Osh4p localization (Alfaro *et al.*, 2011). We speculated that the polarized localization of Osh4p might also be dependent on binding to PI4P based on PI4P's cellular distribution. To test this hypothesis, we first determined if the localization of Osh4 is affected by the loss of enzymatic activities that synthesize PI4P. In budding yeast, there are two major PI 4-kinases that are responsible for PI4P synthesis.

Pik1 is a Golgi-localized lipid kinase while Stt4 functions at the plasma membrane (Audhya *et al.*, 2000). Each of these two PI 4-kinases synthesizes roughly 50% of the total PI4P in cells. We examined the localization of Osh4-3xGFP in *pik1^{ts}* cells and *stt4^{ts}* cells respectively. At the permissive temperature, Osh4-3xGFP localized normally in both *pik1^{ts}* cells and *stt4^{ts}* cells. Interestingly, upon shifting to the non-permissive temperature for 1 hour, Osh4-3xGFP appeared predominantly cytoplasmic in *pik1^{ts}* cells, while its localization was normal in *stt4^{ts}* cells (Figure 3A and Figure 3B). Subcellular fractionation results further support the proposal that Pik1p function is important for Osh4p membrane association (Supplemental Figure 2A and 2B). Together, these results suggest that the localization of Osh4p is controlled by the Pik1-generated pool of PI4P.

To confirm that PI4P binding is required for normal Osh4p localization, we generated two *osh4* alleles harboring point mutations within the PI4P binding pocket, K336A and R344A. Both mutants display defects in PI4P binding but not sterol binding *in vitro* (de Saint-Jean *et al.*, 2011). Consistent with our expectation, neither Osh4 K336A nor R344A mutants were able to localize to polarized secretory sites in cells (Figure 3C and 3D). Western blot analysis revealed that both Osh4p mutant proteins express normally (Figure 3E); thus the microscopy experimental results were not due to the lack of stability of these proteins *in vivo*. Interestingly, the Osh4p mutants defective in PI4P binding were not able to complement the growth of *osh1-7Δ osh4^{ts}* cells at the non-permissive temperature (Figure 3F), indicating the ability to bind PI4P is essential for Osh4 function.

To test whether Osh4p changes localization in response to change of PI4P distribution in cells, we examined the localization of Osh4-3xGFP in wild type cells, *sac1Δ* cells and *inp52Δ inp53Δ* cells, which lack the different phosphatase activities that can dephosphorylate PI4P (Guo *et al.*, 1999a; Hughes *et al.*, 2000; Foti *et al.*, 2001). There was a dramatic increase in the PM localized PI4P pool in *sac1Δ* cells and an increase in the localization of the PI4P probe to the cytoplasmic punctate structures in the *inp52Δ inp53Δ* cells (Figure 4A). Interestingly, we did not see Osh4p shift to the plasma membrane in *sac1Δ* cells. However, we did observe a dramatic increase of large Osh4p-positive puncta in the cytoplasm of *sac1Δ* cells and a significant increase of small Osh4p-positive puncta in the cytoplasm of *inp52Δ inp53Δ* cells (Supplemental Figure 4), suggesting Osh4p's localization was responding to the changes of PI4P distribution distinctly in *sac1Δ* cells and *inp52Δ inp53Δ* cells. In total, our results indicate that PI4P, possibly in concert with additional factors, recruits Osh4p to secretory vesicles.

PI4P accumulates on Sec4p positive vesicles in *osh4Δ* cells

Previous studies, as well as those presented here, support the proposal that PI4P is present on secretory vesicles, at least when they are initially formed (Santiago-Tirado *et al.*, 2011). Furthermore, our data indicate that Osh4 is recruited to secretory vesicles by PI4P. Once it has been recruited, Osh4p could act to reduce the PI4P pool present on secretory vesicles. To test whether loss of Osh4p alone can affect the PI4P distribution in yeast cells, we examined the localization of GFP-2xPH^{Osh2} in wild type and *osh4Δ* cells. In wild type cells, the

GFP-2xPH^{Osh2} was present both on the plasma membrane and Golgi apparatus, as previously reported (Roy and Levine, 2004). Strikingly, we observed in the *osh4Δ* cells a dramatic shift in the PI4P probe to the budding daughter cells that have high concentrations of Sec4p, in comparison to wild type cells (Figure 4A and Figure 4B). Similar results were observed when we assessed the localization of the PH^{FAPP1} PI4P probe. In *osh4Δ* cells, PH^{FAPP1} shifted to structures that are positive for the secretory vesicle marker Sec4p, but not the late Golgi marker Sec7p in the budding daughter cells (Supplemental Figure 5A and 5B). In contrast, we did not observe any localization change for the PI(4,5)P₂ probe, GFP-2xPH^{PLC}, in *osh4Δ* cells (Supplemental Figure 5C). These results suggest that Osh4p plays a role in facilitating either the dephosphorylation or redistribution of PI4P from the secretory vesicles before they become concentrated at exocytic sites.

To better understand how Osh4 affects intracellular PI4P distribution, we captured time-lapse movies of the GFP-2xPH^{Osh2} probe in wild type and *osh4Δ* cells. We observed that the GFP-2xPH^{Osh2} probe was concentrated in the tips of small buds of both wild type and *osh4Δ* cells. However, as daughter cells grew bigger, the polarized localization of the GFP-2xPH^{Osh2} probe was gradually lost in wild type cells (Figure 4C and Movie 1). Interestingly, this process was delayed in the *osh4Δ* cells as we can still observe large GFP-2xPH^{Osh2} positive structures as the cells grew (Figure 4A, Figure 4C and Movie 2). This result suggests that high levels of PI4P accumulate on secretory vesicles in *osh4Δ* cells and, as a

result, the high level of PI4P may inhibit the docking or fusion of these vesicles to the plasma membrane.

The localization patterns of Ypt32p and Sec15p are altered in *osh4Δ* cells

PI4P inhibits the interaction of Sec15p with Sec2p and thereby promotes the interaction of Sec2p with the competing ligand, Ypt32p (Mizuno-Yamasaki *et al.*, 2010). We carried out fluorescence microscopy experiments to ascertain whether Osh4p functions as an upstream factor regulating the localization of Ypt32p and Sec15p perhaps through its effects on the PI4P levels of secretory vesicles. First, we examined the localization of GFP-Ypt32 in wild type and *osh4Δ* cells. As a control, we also include the *sac1Δ* cells and *inp52Δ inp53Δ* cells that are defective in PI4P turnover and may have increased PI4P levels on secretory vesicles. In wild type cells, the majority of Ypt32p localized to the Golgi with a pool of Ypt32p observed at the tips of small budded cells, however, it was rarely observed at polarized secretory sites in large budded cells (Figure 5A and Figure 5C). Strikingly, we observed a dramatic increase of GFP-Ypt32 at polarized secretory sites in large buds of *osh4Δ* cells (Figure 5A). Similar results were observed in the *sac1Δ* cells and *inp52Δ inp53Δ* cells (Figure 5A), suggesting that Osh4p may regulate the localization of Ypt32p by controlling PI4P turnover on secretory vesicles.

Previous results revealed that elevated levels of PI4P inhibit the interaction of Sec15p with Sec2p (Mizuno-Yamasaki *et al.*, 2010) and *osh1-7Δ osh4^{ts}* cells, which have increased levels of PI4P, exhibit Sec15p mislocalization (Supplemental Figure 1B). To test whether loss of Osh4p function or PI4P

phosphatase activity affects Sec15p localization *in vivo*, we examined the localization of Sec15-3xGFP in wild type cells, *osh4Δ* cells, *sac1Δ* cells and *inp52Δ inp53Δ* cells. Unlike the *osh1-7Δ osh4^{ts}* cells, we can still observe Sec15-3xGFP both at small budding tips and mother-daughter cell necks in *osh4Δ* cells (Figure 5B). However, after careful quantification we observed a significant decrease of Sec15p at polarized secretory sites in *osh4Δ* cells, *sac1Δ* cells and *inp52Δ inp53Δ* cells compared to wild type cells (Figure 5C). These results are consistent with the idea that PI4P helps to retain Ypt32p on secretory vesicles while inhibiting the recruitment of Sec15p and subsequent exocyst assembly on secretory vesicles.

Osh4p regulates the Sec2p-Sec15p interaction through PI4P

PI4P inhibits the interaction between Sec2p and Sec15p both *in vivo* and *in vitro* (Mizuno-Yamasaki et al., 2010), therefore we investigated whether this interaction is also controlled by Osh4p *in vivo*. We tagged Sec2p with 3xGFP and Sec15p with 13myc in wild type cells, *osh4Δ* cells and *sac1Δ* cells. We performed coimmuno-precipitation experiments, pulling down Sec2-3xGFP with the anti-GFP antibody and then probing for Sec15p with anti-myc antibody. Consistent with the microscopy data, we observed that in both *osh4Δ* cells and *sac1Δ* cells, less Sec15-13myc was co-immunoprecipitated compared to wild type cells (Figure 6A). Quantification of the co-precipitation experiment revealed that the interaction between Sec2p and Sec15p was roughly 3 fold lower on average in *osh4Δ* cells and *sac1Δ* cells compared to wild type cells (Figure 6B). These data further support the model that Osh4p negatively regulates the PI4P

levels of secretory vesicles and the high levels of PI4P in *osh4* Δ cells inhibits the interaction between Sec2p and Sec15p.

In total, we propose that Osh4p is the principle Osh protein controlling the PI4P level of secretory vesicles. As the secretory vesicles move from Golgi to the plasma membrane, PI4P is turned over and/or extracted under the control of Osh4p (Figure 7). The decrease in the level of PI4P may facilitate the release of Ypt32p and recruitment of Sec15p to secretory vesicles. This mechanism is important for the subsequent assembly of exocyst and tethering of secretory vesicles in preparation for fusion to the plasma membrane.

Discussion

The Golgi and plasma membrane represent the two major sites of PI4P synthesis. Several studies, including this one, indicate that PI4P can also be found in Golgi-derived secretory vesicles and, although these vesicles go on to fuse with the plasma membrane, genetic findings imply that the Golgi pool of PI4P is not efficiently delivered to the plasma membrane (Audhya *et al.*, 2000; Santiago-Tirado *et al.*, 2011). Both the Golgi PI 4-kinase, Pik1p, and the plasma membrane PI 4-kinase, Stt4p, are essential for viability and overexpression of Pik1p fails to complement *stt4* mutants (Audhya *et al.*, 2000). This raises the question of the fate of the PI4P that is incorporated into secretory vesicles as they bud from the Golgi. The findings presented here support the proposal that the vesicular pool of PI4P is normally reduced prior to vesicle fusion with the plasma membrane and that this process involves the oxysterol binding protein, Osh4p (Figure 7). Dephosphorylation by the PI4P phosphatase, Sac1p as well as

possibly, Inp52p and Inp53p, proteins that contain Sac1 lipid phosphatase domains, appears to be a likely mechanism, although we must also consider alternative mechanisms.

Our results show that Osh4p is recruited to secretory vesicles in a PI4P-dependent manner. The Osh4p K336A and R344A mutants are defective in PI4P binding but not in ergosterol binding *in vitro* (de Saint-Jean *et al.*, 2011) and they exhibit a loss of localization *in vivo*. Therefore, the association of Osh4p with specific membranes is likely regulated by the local PI4P concentration. However, PI4P is probably not sufficient for Osh4p recruitment since in a *sac1Δ* mutant strain the level of PI4P on the plasma membrane is greatly increased without a concomitant shift in Osh4p localization to the cell cortex. The membrane curvature-sensing ALPS domain of Osh4p (Drin *et al.*, 2007) could play an important role in the recruitment of Osh4p to 100nm secretory vesicles in preference to the relatively flat plasma membrane. Sec4p is also required for Osh4p recruitment, but apparently not via a direct interaction.

Osh proteins have been shown to stimulate Sac1p phosphatase activity *in vitro* and loss of either Osh protein function or Sac1p function leads to a dramatic increase in PI4P levels *in vivo* (Foti *et al.*, 2001; Stefan *et al.*, 2011). Nonetheless, Sac1p is an integral membrane protein found predominantly on the ER membrane with a smaller pool associated with the Golgi and there is no evidence that Sac1p is incorporated into Golgi-derived secretory vesicles (Faulhammer *et al.*, 2007). This raises the question of how Sac1p on the ER can impact PI4P levels on secretory vesicles. Osh4p can extract PI4P from one set of

liposomes and exchange it for ergosterol in another set of liposomes *in vitro* (de Saint-Jean et al., 2011). This observation has prompted the hypothesis that Osh proteins, such as Osh4p, could extract PI4P from cellular membranes such as secretory vesicles and present it to enzymes that metabolize it on another membrane, such as Sac1p on the ER. Another possibility is that Osh proteins could present PI4P to the phosphatase at organelle contact sites. The plasma membrane bound pool of PI4P is de-phosphorylated by Sac1p at ER-PM contact sites in a reaction that requires Osh proteins (Stefan *et al.*, 2011; Manford *et al.*, 2012). Osh4p on secretory vesicles could present PI4P to the Sac1p phosphatase *in trans* as vesicles pass alongside the ER membrane on their way to exocytic sites. Although secretory vesicles do not bind in a stable fashion to the ER, most of the cell cortex is lined with a mix of ER sheets and tubules so vesicles could transiently associate with the cortical ER prior to fusion with the plasma membrane. The PI4P on secretory vesicles might also be presented by Osh4p to the PI4P 5-kinase, Mss4, at the plasma membrane and be converted to PI(4,5)P₂ if the vesicles were in proximity to the Mss4 PIK patches (Ling *et al.*, 2012). This could explain why the PI(4,5)P₂ level is surprisingly low in *osh1-7Δ osh4^{ts}* cells at the nonpermissive temperature even though the PI4P level is highly elevated (Stefan *et al.*, 2011). Further studies on the Osh proteins will be needed to clarify these important questions.

The Osh4p-dependent reduction in the level of PI4P on secretory vesicles represents an important vesicle maturation event that helps to direct a switch in the binding partners of Sec2p from an initial interaction with Ypt31/32-GTP to a

competing interaction with Sec15p (Figure 7) (Mizuno-Yamasaki *et al.*, 2010). Supporting this model, in *osh4* or *sac1* mutants there is a decrease in the coprecipitation of Sec15p with Sec2p and a shift is seen in the distribution of Ypt32p towards exocytic sites and in the distribution of Sec15p away from exocytic sites. This switch in Sec2p's binding partners is also under regulation by phosphorylation (Stalder *et al.*, 2013). Sec2p is phosphorylated within the region responsible for binding both Ypt31/32-GTP and Sec15p and the phosphorylated form has a strong preference for binding Sec15p while the non-phosphorylated form preferentially binds Ypt32-GTP (Stalder *et al.*, 2013). These two vesicle maturation events, PI4P reduction and Sec2p phosphorylation, may help to prepare the vesicles for tethering and fusion to the plasma membrane.

Materials and Methods

Yeast strains and Plasmid Construction

A list of *Saccharomyces cerevisiae* strains and plasmids used in this study and their genotypes can be found in Supplementary Tables S1 and S2. Homologous recombination was used to tag or delete genes in yeast (Longtine *et al.*, 1998). All integrations and deletions were verified by PCR analysis and expression of fusion proteins was confirmed by Western Blot analysis. The yeast shuttle vectors used in this study have been previously described (Sikorski and Hieter, 1989).

Fluorescence Microscopy and Analysis of Images

Yeast cells were grown overnight to mid-log phase (OD₆₀₀ of 0.3-0.5) in a synthetic medium containing 2% glucose at 25 °C. For temperature-sensitive strains, the cell culture was shifted to 37 °C for 60 minutes or as described.

Fluorescence images were acquired with a 63x or 100x oil immersion objective (Carl Zeiss) on a spinning disc confocal microscopy system (Yokagawa Corporation of America), which included a microscope (Observer Z1; Carl Zeiss) equipped with an electron multiplying CCD camera (QuantEM 512SC; Photometrics). Excitation of GFP, mCherry or DsRed was achieved using 488-nm argon and 568-nm argon/krypton lasers. For most of the samples, a z-stack of 12 slices with a 270-nm slice distance was generated. For samples in which the localizations of two proteins were examined, only images from one z section are shown. Images were analyzed using AxioVision software 4.8 (Carl Zeiss). Levels of the fluorescence images and DIC images were adjusted with Adobe Photoshop, the same parameters were applied to all images within the same figure. Osh4-3xGFP, GFP-Ypt32 and Sec15-3xGFP localization were scored based on localization at the tips of small buds or the tips/necks of large buds. 100–200 cells were examined for each condition and three independent experiments were used to calculate the SD.

Label Plasma Membrane with FM4-64

5 OD₆₀₀ equivalents of mid-log cells were harvested and re-suspended in 100 μ L ddH₂O on ice. 5 μ L cells were mixed with 5 μ L FM4-64 solution (20 μ g/mL FM4-64, 20 mM Tris-HCl pH 7.5, 20 mM NaN₃, 20 mM NaF) on ice. Cells were immediately used for microscopy experiments.

Analysis of Cellular Protein Expression Levels

5 OD₆₀₀ equivalents of mid-log cells were harvested by precipitation in 10% trichloroacetic acid (TCA). Precipitates were washed in acetone, aspirated,

resuspended in lysis buffer (150 mM NaCl, 50mM Tris pH 7.5, 1mM EDTA, 1% SDS), and mechanically lysed with silica beads. Protein sample buffer (150 mM Tris pH 6.8, 6M Urea, 6%SDS, 10% β -mercaptoethanol, 20% Glycerol) was added and extracts were analyzed by SDS-PAGE and immunoblotting with anti-GFP (Santa Cruz Biotechnology) or anti-Adh1 antibodies.

Coimmunoprecipitation Assay

Yeast cells expressing epitope-tagged proteins were grown to an OD₆₀₀ of 0.5-0.8. Cells were harvested and spheroplasted with lyticase (Scott and Schekman, 1980). Spheroplasts were lysed mechanically in the lysis buffer (1xPBS, 5mM EDTA, pH=8.0 with protease inhibitors added) by using a homogenizer (Wheaton). Cell lysates were treated with 0.5% Tween-20 and cleared by centrifugation at 13,000x g for 10 minutes. Immunoprecipitations were performed by adding anti-GFP antibody (Ferro-Novick lab) to the supernatant at 4°C and incubated over night. The protein A/G resin (Thermo Scientific) were added to the samples and further incubated for 2 hours at 4°C next morning. After washing the affinity resin with PBS IP buffer four times, immunoprecipitated proteins were eluted and reduced with sample buffer (6M Urea, 150mM Tris-HCl pH=6.8, 6% SDS, 10% β -mercaptoethanol, Bromophenol Blue) and analyzed by SDS-PAGE and immunoblotting. The intensities of western-blot bands were analyzed using NIH ImageJ software.

Acknowledgements

We thank S. Ferro-Novick for anti-GFP antibody and B. Antony and G. Drin for the *osh4* K336A and R344A mutants. We thank S. Field for the GFP-2xPH(PLC1delta) plasmid. We thank Y. Jones from the lab of M. Farquhar for preparation of samples for electron microscopy. This study was supported by grants GM082861 and GM35370 from the N.I.H. to P.N.

References:

- Alfaro, G., Johansen, J., Dighe, S.A., Duamel, G., Kozminski, K.G., and Beh, C.T. (2011). The sterol-binding protein Kes1/Osh4p is a regulator of polarized exocytosis. *Traffic* 12, 1521-1536.
- Audhya, A., and Emr, S.D. (2002). Stt4 PI 4-kinase localizes to the plasma membrane and functions in the Pkc1-mediated MAP kinase cascade. *Dev Cell* 2, 593-605.
- Audhya, A., Foti, M., and Emr, S.D. (2000). Distinct roles for the yeast phosphatidylinositol 4-kinases, Stt4p and Pik1p, in secretion, cell growth, and organelle membrane dynamics. *Mol Biol Cell* 11, 2673-2689.
- Bacon, R.A., Salminen, A., Ruohola, H., Novick, P., and Ferro-Novick, S. (1989). The GTP-binding protein Ypt1 is required for transport in vitro: the Golgi apparatus is defective in *ypt1* mutants. *J Cell Biol* 109, 1015-1022.
- Balla, T. (2013). Phosphoinositides: tiny lipids with giant impact on cell regulation. *Physiol Rev* 93, 1019-1137.
- Beh, C.T., Cool, L., Phillips, J., and Rine, J. (2001). Overlapping functions of the yeast oxysterol-binding protein homologues. *Genetics* 157, 1117-1140.
- Beh, C.T., and Rine, J. (2004). A role for yeast oxysterol-binding protein homologs in endocytosis and in the maintenance of intracellular sterol-lipid distribution. *J Cell Sci* 117, 2983-2996.
- Behnia, R., and Munro, S. (2005). Organelle identity and the signposts for membrane traffic. *Nature* 438, 597-604.

- Benli, M., Doring, F., Robinson, D.G., Yang, X., and Gallwitz, D. (1996). Two GTPase isoforms, Ypt31p and Ypt32p, are essential for Golgi function in yeast. *EMBO J* 15, 6460-6475.
- Carlton, J.G., and Cullen, P.J. (2005). Coincidence detection in phosphoinositide signaling. *Trends in cell biology* 15, 540-547.
- Chen, S., Novick, P., and Ferro-Novick, S. (2012). ER network formation requires a balance of the dynamin-like GTPase Sey1p and the Lunapark family member Lnp1p. *Nat Cell Biol* 14, 707-716.
- de Saint-Jean, M., Delfosse, V., Douguet, D., Chicanne, G., Payraastre, B., Bourguet, W., Antonny, B., and Drin, G. (2011). Osh4p exchanges sterols for phosphatidylinositol 4-phosphate between lipid bilayers. *J Cell Biol* 195, 965-978.
- Donovan, K.W., and Bretscher, A. (2012). Myosin-V is activated by binding secretory cargo and released in coordination with Rab/exocyst function. *Dev Cell* 23, 769-781.
- Drin, G., Casella, J.F., Gautier, R., Boehmer, T., Schwartz, T.U., and Antonny, B. (2007). A general amphipathic alpha-helical motif for sensing membrane curvature. *Nat Struct Mol Biol* 14, 138-146.
- Faulhammer, F., Kanjilal-Kolar, S., Knodler, A., Lo, J., Lee, Y., Konrad, G., and Mayinger, P. (2007). Growth control of Golgi phosphoinositides by reciprocal localization of sac1 lipid phosphatase and pik1 4-kinase. *Traffic* 8, 1554-1567.
- Foti, M., Audhya, A., and Emr, S.D. (2001). Sac1 lipid phosphatase and Stt4 phosphatidylinositol 4-kinase regulate a pool of phosphatidylinositol 4-phosphate that functions in the control of the actin cytoskeleton and vacuole morphology. *Mol Biol Cell* 12, 2396-2411.
- Govindan, B., Bowser, R., and Novick, P. (1995). The role of Myo2, a yeast class V myosin, in vesicular transport. *J Cell Biol* 128, 1055-1068.
- Grosshans, B.L., Andreeva, A., Gangar, A., Niessen, S., Yates, J.R., 3rd, Brennwald, P., and Novick, P. (2006). The yeast Igl family member Sro7p is an effector of the secretory Rab GTPase Sec4p. *J Cell Biol* 172, 55-66.
- Guo, S., Stolz, L.E., Lemrow, S.M., and York, J.D. (1999a). SAC1-like domains of yeast SAC1, INP52, and INP53 and of human synaptojanin encode polyphosphoinositide phosphatases. *J Biol Chem* 274, 12990-12995.
- Guo, W., Roth, D., Walch-Solimena, C., and Novick, P. (1999b). The exocyst is an effector for Sec4p, targeting secretory vesicles to sites of exocytosis. *EMBO J* 18, 1071-1080.

Hughes, W.E., Woscholski, R., Cooke, F.T., Patrick, R.S., Dove, S.K., McDonald, N.Q., and Parker, P.J. (2000). SAC1 encodes a regulated lipid phosphoinositide phosphatase, defects in which can be suppressed by the homologous Inp52p and Inp53p phosphatases. *J Biol Chem* 275, 801-808.

Hutagalung, A.H., and Novick, P.J. (2011). Role of Rab GTPases in membrane traffic and cell physiology. *Physiol Rev* 91, 119-149.

Im, Y.J., Raychaudhuri, S., Prinz, W.A., and Hurley, J.H. (2005). Structural mechanism for sterol sensing and transport by OSBP-related proteins. *Nature* 437, 154-158.

Jedd, G., Mulholland, J., and Segev, N. (1997). Two new Ypt GTPases are required for exit from the yeast trans-Golgi compartment. *J Cell Biol* 137, 563-580.

Jedd, G., Richardson, C., Litt, R., and Segev, N. (1995). The Ypt1 GTPase is essential for the first two steps of the yeast secretory pathway. *J Cell Biol* 131, 583-590.

Jin, Y., Sultana, A., Gandhi, P., Franklin, E., Hamamoto, S., Khan, A.R., Munson, M., Schekman, R., and Weisman, L.S. (2011). Myosin V transports secretory vesicles via a Rab GTPase cascade and interaction with the exocyst complex. *Dev Cell* 21, 1156-1170.

Kozminski, K.G., Alfaro, G., Dighe, S., and Beh, C.T. (2006). Homologues of oxysterol-binding proteins affect Cdc42p- and Rho1p-mediated cell polarization in *Saccharomyces cerevisiae*. *Traffic* 7, 1224-1242.

LeBlanc, M.A., and McMaster, C.R. (2010). Lipid binding requirements for oxysterol-binding protein Kes1 inhibition of autophagy and endosome-trans-Golgi trafficking pathways. *J Biol Chem* 285, 33875-33884.

Levine, T.P., and Munro, S. (2001). Dual targeting of Osh1p, a yeast homologue of oxysterol-binding protein, to both the Golgi and the nucleus-vacuole junction. *Mol Biol Cell* 12, 1633-1644.

Ling, Y., Stefan, C.J., Macgurn, J.A., Audhya, A., and Emr, S.D. (2012). The dual PH domain protein Opy1 functions as a sensor and modulator of PtdIns(4,5)P(2) synthesis. *EMBO J* 31, 2882-2894.

Longtine, M.S., McKenzie, A., 3rd, Demarini, D.J., Shah, N.G., Wach, A., Brachat, A., Philippsen, P., and Pringle, J.R. (1998). Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* 14, 953-961.

- Manford, A.G., Stefan, C.J., Yuan, H.L., Macgurn, J.A., and Emr, S.D. (2012). ER-to-plasma membrane tethering proteins regulate cell signaling and ER morphology. *Dev Cell* 23, 1129-1140.
- McNew, J.A., Parlati, F., Fukuda, R., Johnston, R.J., Paz, K., Paumet, F., Sollner, T.H., and Rothman, J.E. (2000). Compartmental specificity of cellular membrane fusion encoded in SNARE proteins. *Nature* 407, 153-159.
- Medkova, M., France, Y.E., Coleman, J., and Novick, P. (2006). The rab exchange factor Sec2p reversibly associates with the exocyst. *Mol Biol Cell* 17, 2757-2769.
- Mizuno-Yamasaki, E., Medkova, M., Coleman, J., and Novick, P. (2010). Phosphatidylinositol 4-phosphate controls both membrane recruitment and a regulatory switch of the Rab GEF Sec2p. *Dev Cell* 18, 828-840.
- Mizuno-Yamasaki, E., Rivera-Molina, F., and Novick, P. (2012). GTPase networks in membrane traffic. *Annu Rev Biochem* 81, 637-659.
- Ortiz, D., Medkova, M., Walch-Solimena, C., and Novick, P. (2002). Ypt32 recruits the Sec4p guanine nucleotide exchange factor, Sec2p, to secretory vesicles; evidence for a Rab cascade in yeast. *J Cell Biol* 157, 1005-1015.
- Roy, A., and Levine, T.P. (2004). Multiple pools of phosphatidylinositol 4-phosphate detected using the pleckstrin homology domain of Osh2p. *J Biol Chem* 279, 44683-44689.
- Santiago-Tirado, F.H., Legesse-Miller, A., Schott, D., and Bretscher, A. (2011). PI4P and Rab inputs collaborate in myosin-V-dependent transport of secretory compartments in yeast. *Dev Cell* 20, 47-59.
- Schott, D., Ho, J., Pruyne, D., and Bretscher, A. (1999). The COOH-terminal domain of Myo2p, a yeast myosin V, has a direct role in secretory vesicle targeting. *J Cell Biol* 147, 791-808.
- Schott, D.H., Collins, R.N., and Bretscher, A. (2002). Secretory vesicle transport velocity in living cells depends on the myosin-V lever arm length. *J Cell Biol* 156, 35-39.
- Schulz, T.A., Choi, M.G., Raychaudhuri, S., Mears, J.A., Ghirlando, R., Hinshaw, J.E., and Prinz, W.A. (2009). Lipid-regulated sterol transfer between closely apposed membranes by oxysterol-binding protein homologues. *J Cell Biol* 187, 889-903.
- Sclafani, A., Chen, S., Rivera-Molina, F., Reinisch, K., Novick, P., and Ferro-Novick, S. (2010). Establishing a role for the GTPase Ypt1p at the late Golgi. *Traffic* 11, 520-532.

Scott, J.H., and Schekman, R. (1980). Lyticase: endoglucanase and protease activities that act together in yeast cell lysis. *J Bacteriol* *142*, 414-423.

Shen, D., Yuan, H., Hutagalung, A., Verma, A., Kummel, D., Wu, X., Reinisch, K., McNew, J.A., and Novick, P. (2013). The synaptobrevin homologue Snc2p recruits the exocyst to secretory vesicles by binding to Sec6p. *J Cell Biol* *202*, 509-526.

Sikorski, R.S., and Hieter, P. (1989). A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* *122*, 19-27.

Stalder, D., Mizuno-Yamasaki, E., Ghassemian, M., and Novick, P.J. (2013). Phosphorylation of the Rab exchange factor Sec2p directs a switch in regulatory binding partners. *Proc Natl Acad Sci U S A* *110*, 19995-20002.

Stefan, C.J., Manford, A.G., Baird, D., Yamada-Hanff, J., Mao, Y., and Emr, S.D. (2011). Osh proteins regulate phosphoinositide metabolism at ER-plasma membrane contact sites. *Cell* *144*, 389-401.

TerBush, D.R., Maurice, T., Roth, D., and Novick, P. (1996). The Exocyst is a multiprotein complex required for exocytosis in *Saccharomyces cerevisiae*. *EMBO J* *15*, 6483-6494.

Wagner, W., Bielli, P., Wacha, S., and Ragnini-Wilson, A. (2002). Mlc1p promotes septum closure during cytokinesis via the IQ motifs of the vesicle motor Myo2p. *EMBO J* *21*, 6397-6408.

Walch-Solimena, C., Collins, R.N., and Novick, P.J. (1997). Sec2p mediates nucleotide exchange on Sec4p and is involved in polarized delivery of post-Golgi vesicles. *J Cell Biol* *137*, 1495-1509.

Walch-Solimena, C., and Novick, P. (1999). The yeast phosphatidylinositol-4-OH kinase pik1 regulates secretion at the Golgi. *Nat Cell Biol* *1*, 523-525.

Table 1. Yeast strains used in this study.

Strain	Genotype	Reference
NY1211	MAT α <i>his3-Δ200 leu2-3,112 ura3-52 GAL⁺</i>	Lab strain
CBY924	MAT α <i>ura3-52 his3-Δ200 leu2-3, 112 trp1-Δ901 suc2-Δ9 lys2-801 osh1Δ::kan-MX4 osh2Δ::kan-MX4 osh3Δ::LYS2 osh4Δ::HIS3 osh5Δ::LEU2\square osh6Δ::LEU2 osh7Δ::HIS / OSH4, CEN-TRP</i>	Beh CT and Rine J, 2004
CBY926	MAT α <i>ura3-52 his3-Δ200 leu2-3, 112 trp1-Δ901 suc2-Δ9 lys2-801 osh1Δ::kan-MX4 osh2 Δ::kan-MX4 osh3Δ::LYS2 osh4Δ::HIS3 osh5Δ::LEU2\square osh6Δ::LEU2 osh7Δ::HIS3 / osh4-1, CEN-TRP</i>	Beh CT and Rine J, 2004
NY3076	NY1211 <i>osh4::[OSH4-3GFP HIS3]</i>	This study
NY3077	NY1211 <i>osh4::[OSH4-3GFP HIS3] leu2-3,112::[mcherry-SEC4 LEU2]</i>	This study
NY408	MAT α <i>ura3-52 his4-619 sec4-8</i>	Lab strain
NY3078	NY1211 <i>osh4::[OSH4-3GFP URA3]</i>	This study
NY3079	NY408 <i>osh4::[OSH4-3GFP URA3]</i>	This study
AAY102	MAT α <i>ura3-52 his3-Δ200 leu2-3,112 trp1-Δ901 suc2-Δ9 lys2-801 stt4Δ::HIS3 pRS415 stt4-4</i>	Audhya et al., 2000
NY2189	MAT α <i>leu2-3,112 ura3-52 pik1-101 GAL⁺</i>	Lab strain
NY3080	AAY102 <i>osh4::[OSH4-3GFP URA3]</i>	This study
NY3081	NY2189 <i>osh4::[OSH4-3GFP URA3]</i>	This study
BY4741	MAT α <i>his3Δ1 leu2Δ0 met15Δ0 ura3Δ0</i>	Lab strain
NY3082	BY4741 <i>inp52Δ::HIS3 inp53Δ</i>	This study
NY3083	BY4741 <i>ura3-52::[GFP-YPT32 URA3]</i>	This study
NY3084	BY4741 <i>sac1Δ ura3-52::[GFP-YPT32 URA3]</i>	This study
NY3085	BY4741 <i>osh4Δ ura3-52::[GFP-YPT32 URA3]</i>	This study
NY3086	BY4741 <i>inp52Δ::HIS3 inp53Δ ura3-52::[GFP-YPT32 URA3]</i>	This study
NY3087	BY4741 <i>SEC15::[SEC15-3GFP URA3]</i>	This study
NY3088	BY4741 <i>sac1Δ SEC15::[SEC15-3GFP URA3]</i>	This study
NY3089	BY4741 <i>osh4Δ SEC15::[SEC15-3GFP URA3]</i>	This study
NY3090	BY4741 <i>inp52Δ::HIS3 inp53Δ SEC15::[SEC15-3GFP URA3]</i>	This study
NY3091	BY4741 <i>SEC15::[SEC15-13myc URA3]</i>	This study
NY3092	BY4741 <i>sac1Δ SEC15::[SEC15-13myc URA3]</i>	This study
NY3093	BY4741 <i>osh4Δ SEC15::[SEC15-13myc URA3]</i>	This study
NY3094	BY4741 <i>SEC15::[SEC15-13myc URA3] sec2::[SEC2-3GFP LEU2]</i>	This study
NY3095	BY4741 <i>sac1Δ SEC15::[SEC15-13myc URA3] sec2::[SEC2-3GFP LEU2]</i>	This study
NY3096	BY4741 <i>osh4Δ SEC15::[SEC15-13myc URA3] sec2::[SEC2-3GFP LEU2]</i>	This study
NY3097	NY1211 <i>osh4::[osh4-3GFP K336A URA3]</i>	This study
NY3098	NY1211 <i>osh4::[osh4-3GFP R344A URA3]</i>	This study

Table 2. Plasmids used in this study.

Plasmid	Description	Source/Reference
NRB1445	<i>pRS306-pADH-GFP-FAPP1-PH</i>	Lab strain
NRB1444	<i>pRS306-pADH-mCherry-FAPP1-PH</i>	Lab strain
SFNB797	<i>pUSE-URA3-SEC7-EGFP</i>	Susan Ferro-Novick Lab
NRB1584	<i>pRS303-OSH4(bp 730-1302)-3GFP-tADH</i>	This study
NRB1585	<i>pRS306-OSH4(bp 730-1302)-3GFP-tADH</i>	This study
NRB139	<i>YCp50-URA3-SEC4</i>	Lab strain, Walworth NC et al. 1989
NRB170	<i>Ycp50-URA3-SEC4 (Overexpression)</i>	Lab strain, Walworth NC et al. 1989
pTL511	<i>pRS416-2x(GFP-Osh2-PH)</i>	Roy A and Levine TP, 2004
NRB1347	<i>Yiplac128-pADH-mCherry-SEC4-tCYC1</i>	Lab strain
NRB1325	<i>pRS306-pADH-GFP-YPT32-tCYC1</i>	Lab strain
NRB1308	<i>pRS306-SEC15-3GFP</i>	Lab strain
NRB1586	<i>pRS306-SEC15-13myc</i>	This study
NRB1416	<i>pRS305-SEC2(bp 122-2277)-3GFP-tADH</i>	Lab strain
NRB1587	<i>pRS306-OSH4(bp 730-1302) K336A-3GFP-tADH</i>	This study
NRB1588	<i>pRS306-OSH4(bp 730-1302) R344A-3GFP-tADH</i>	This study
NRB1589	<i>pET27-10xHis-OSH4</i>	This study
NRB1590	<i>pGEX4T1-GST-SEC4</i>	This study
NRB1591	<i>pGEX4T1-GST-sec4(S34N)</i>	This study
NRB1592	<i>pGEX4T1-GST-sec4(Q79L)</i>	This study
NRB1440	<i>pGEX4T1-GST-YPT32</i>	Lab strain
NRB1593	<i>pRS316-pOSH4-OSH4</i>	This study
NRB1594	<i>pRS316-pOSH4-OSH4-3GFP-tADH</i>	This study
NRB1595	<i>pRS316-pOSH4-osh4(K336A)-3GFP-tADH</i>	This study
NRB1596	<i>pRS316-pOSH4-osh4(R344A)-3GFP-tADH</i>	This study
NRB1597	<i>pRS426-GFP-2xPH(PLC1delta)</i>	Seth Field Lab

Ling et al., Figure 1




Figure 1. Osh proteins regulate PI4P distribution in cells

(A) Localization of GFP-PH^{FAPP1} in *osh1-7Δ /CEN OSH4* cells or *osh1-7Δ /CEN osh4^{ts}* cells at indicated temperatures. Cells were grown overnight at 25 °C in a synthetic medium containing 2% glucose and then shifted to 37°C for 1 hour. Cells shown are representative of over 100 cells observed. 6% and 55% of *osh1-7Δ /CEN osh4^{ts}* cells showed polarized localization of GFP-PH^{FAPP1} at 25°C and 37°C, respectively. 6% and 0% of *osh1-7Δ /CEN OSH4* cells showed polarized localization of GFP-PH^{FAPP1} at 25°C and 37°C, respectively. Scale bar, 5 μm.

(B) Localization of GFP-PH^{FAPP1} and FM4-64 in *osh1-7Δ /CEN osh4^{ts}* cells at 37 °C. Cells were grown overnight at 25 °C in a synthetic medium containing 2% glucose and then shifted to 37°C for 1 hour. FM4-64 was added to cells and kept on ice to label the plasma membrane. Scale bar, 5 μm.

(C) The PI4P probe GFP-2xPH^{Osh2} accumulates at polarized growth sites in *sec6^{ts}* cells at the non-permissive temperature. Wild type and *sec6^{ts}* cells were grown overnight at 25 °C in a synthetic medium containing 2% glucose and then shifted to 37°C for 15 minutes. Cells shown are representative of over 100 cells observed. Scale bar, 5 μm.

Ling et al., Figure 2






Figure 2. Osh4 is present on Sec4 positive structures in cells

(A) Localization of mCherry-Sec4 and Osh4-3xGFP in cells. Cells were grown overnight at 25 °C in a synthetic medium containing 2% glucose. The boxed area shows a region magnified four-fold. Arrows indicate Osh4-3xGFP is present on cytoplasmic punctate structures. Arrowheads indicate Osh4-3xGFP colocalize with mCherry-Sec4 in small buds. Scale bar, 2 μ m.

(B) The polarized localization of Osh4-3xGFP is dependent on Sec4. Osh4-3xGFP localization in wild type and *sec4^{ts}* cells at indicated temperatures. Cells were grown overnight at 25 °C in a synthetic medium containing 2% glucose and then shifted to 37°C for 1 hour. Scale bar, 5 μ m.

(C) Quantification of wild type cells and *sec4^{ts}* cells with polarized localized Osh4-3xGFP at 25 °C or 37°C for 1 hour. A total of 200 cells were counted for each experiment. Values indicate the percentage of cells showing bud or mother-daughter neck localization of Osh4-3xGFP. Mean and SD of three experiments are shown.

Ling et al., Figure 3




Figure 3. The polarized localization of Osh4 is dependent on PI4P

(A) Localization of Osh4-3xGFP in *stt4^{ts}* cells and *pik1^{ts}* cells at indicated temperatures. Osh4-3xGFP is mislocalized in *pik1^{ts}* cells at the non-permissive

temperature. Cells were grown overnight at 25 °C in a synthetic medium containing 2% glucose and then shifted to 37°C for 1 hour. Scale bar, 5 μm.

(B) Quantification of *stt4^{ts}* cells and *pik1^{ts}* cells with regards to polarized localization of Osh4-3xGFP at 25 °C or 37°C for 1 hour. A total of 200 cells were counted for each experiment. Values indicate the percentage of cells showing bud or mother-daughter neck localization of Osh4-3xGFP. Mean and SD of three experiments are shown. Note: there were no *pik1^{ts}* cells observed with polarized Osh4-3xGFP at 37°C.

(C) Localization of Osh4-3xGFP point mutants defective in PI4P binding in cells. Scale bar, 2 μm.


(D) Quantification of cells with polarized localized Osh4-3xGFP point mutant proteins. A total of 200 cells were counted for each experiment. Values indicate the percentage of cells showing bud or mother-daughter neck localization of Osh4-3xGFP. Mean and SD of three experiments are shown. Note: there were no large budded cells observed with polarized Osh4 K396A-3xGFP or Osh4 R344A-3xGFP.

(E) Steady-state expression levels of Osh4-3xGFP, Osh4^{K336A}-3xGFP, and Osh4^{R344A}-3xGFP. Yeast whole cell lysates were prepared from strains expressing different Osh4 mutant proteins as described in experimental procedures. Adh1p was followed as a loading control.


(F) Complementation assays of Osh4p point mutants. The *osh1-7Δ /CEN osh4^{ts}* cells were transformed with plasmids expressing various mutant Osh4p proteins as indicated. Serial dilutions of yeast cells were grown on –Ura plates to retain plasmids at 37°C for 3 days; only cells harboring functional *OSH4* plasmids grew on plates.

Ling et al., Figure 4

A



B



C




Figure 4. Osh4 regulates PI4P distribution and dynamics in cells

(A) Localization of the PI4P probe GFP-2xPH^{Osh2} in wild type cells, *osh4Δ* cells, *sac1Δ* cells and *inp52Δ inp53Δ* cells. Cells shown are representative of over 100 cells observed. Scale bar, 5 μm.

(B) Localization of the PI4P probe GFP-2xPH^{Osh2} and mCherry-Sec4 in wild type cells and *osh4Δ* cells. Cells shown are representative of over 100 cells observed. Scale bar, 5 μm.

(C) PI4P dynamics in wild type cells and *osh4Δ* cells. Cells expressing GFP-2xPH^{Osh2} were examined by time-lapse fluorescence microscopy. Images at indicated time points are shown. Arrows indicate GFP-2xPH^{Osh2} were accumulated at polarized growth sites. Scale bar, 2 μm.




Figure 5. Osh4 regulates the distribution of Ypt32 and Sec15 in cells
 (A) Localization of GFP-Ypt32 in wild type cells, *osh4Δ* cells, *sac1Δ* cells and *inp52Δ inp53Δ* cells. Scale bar, 5 μ m.

(B) Localization of Sec15-3xGFP in wild type cells, *osh4* Δ cells, *sac1* Δ cells and *inp52* Δ *inp53* Δ cells. Scale bar, 5 μ m.

(C) Quantification of wild type cells, *osh4* Δ cells, *sac1* Δ cells and *inp52* Δ *inp53* Δ cells with polarized localized GFP-Ypt32 or Sec15-3xGFP. A total of 200 cells were counted for each experiment. Values indicate the percentage of cells showing bud or mother-daughter neck localization of GFP-Ypt32 or Sec15-3xGFP. Mean and SD of three experiments are shown.

Ling et al., Figure 6




Figure 6. Osh4 positively regulates the Sec2-Sec15 interaction

(A) The Sec2-Sec15 interaction is inhibited in *osh4Δ* cells and *sac1Δ* cells. Sec2-3xGFP was immunoprecipitated with anti-GFP antibody from wild type, *osh4Δ* or *sac1Δ* cell lysates. As a negative control, wild type cells expressing non-tagged Sec2 were included in the co-IP experiments. 0.5% of whole cell lysates were loaded as input, the rest of the samples were processed for coimmunoprecipitation. Sec2-3xGFP in the immunoprecipitates was detected with anti-Sec2 antibody. Co-precipitated Sec15-13xmyc was detected with anti-myc antibody.

(B) Quantification of the Sec2-Sec15 interaction in wild type cells, *osh4* Δ cells and *sac1* Δ cells. Three independent experiments were performed. The intensity of the bands was quantified using ImageJ. The percentage of Sec15 in the immunoprecipitate was calculated and indicated. The mean and SD of three experiments are shown.




Figure 7. A model for Osh4's role in regulating PI4P during secretion
 Osh4 is recruited to the secretory vesicles through PI4P binding (and possibly through protein-protein interactions as well). Osh4 negatively regulates PI4P levels on the secretory vesicles and thereby enhances the Sec2-Sec15 interaction during this process. Finally, Osh4 leaves the secretory vesicles once PI4P levels are reduced below a certain threshold.